

Source details

Vestnik Zoologii

Continued as: Zoodiversity

Open Access (i)

Scopus coverage years: from 2009 to 2019, from 2021 to Present

(coverage discontinued in Scopus)

Publisher: Naukova Dumka

ISSN: 0084-5604 E-ISSN: 2073-2333

Subject area: (Agricultural and Biological Sciences: Animal Science and Zoology)

(Agricultural and Biological Sciences: Ecology, Evolution, Behavior and Systematics)

Source type: Journal

View all documents >

Set document alert

■ Save to source list Source Homepage

CiteScore CiteScore rank & trend Scopus content coverage

Improved CiteScore methodology


CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. Learn more >

CiteScore 2019

$$1.0 = \frac{215 \text{ Citations } 2016 - 2019}{215 \text{ Documents } 2016 - 2019}$$

Calculated on 06 May, 2020

CiteScore rank 2019 ①



View CiteScore methodology > CiteScore FAQ > Add CiteScore to your site &

CiteScore 2019

1.0

SJR 2021

0.217

(i)

①

×

Morphology

Vestnik Zoologii, **53**(2): 165–172, 2019 DOI: 10.2478/vzoo-2019-0017



UDC 597.551.2:612.335

HISTOLOGICAL AND ULTRASTRUCTURAL STUDIES ON THE INTESTINE OF GUNTEA LOACH, LEPIDOCEPHALICHTHYS GUNTEA (CYPRINIFORMES, COBITIDAE)

S. K. Ghosh

Department of Zoology, Bejoy Narayan Mahavidyalaya, Itachuna, Hooghly-712 147, West Bengal, India

E-mail: saroj.fisherylab@gmail.com

Histological and Ultrastructural Studies on the Intestine of Guntea loach, Lepidocephalichthys guntea (Cypriniformes Cobitidae). Ghosh, S. K. — The cellular organizations of intestine in Lepidocephalichthys guntea (Hamilton, 1822) have been described by light as well as scanning and transmission electron microscopy. The intestine is short and straight like, marked into anterior, middle and posterior region based on mucosal folds, number and size of columnar epithelial cells and mucous cells, thickness of submucosa and muscularis layer. The mucosa of anterior intestine forms high folds, which are lined with compactly arranged columnar epithelial cells and mucous cells. In the middle intestine, folds are pointless whereas the posterior intestine is without folds. The submucosa is formed of thin layer of connective tissue, contained collagen bundles and blood capillaries, comparatively well developed in the posterior intestine. By scanning electron microscopy, outlines of the luminal surface of anterior and middle intestine is embossed with oval or rounded columnar epithelial cells contained densely packed stubby microridges. The posterior intestine has closely set longitudinal folds characterized with minute blood capillaries and columnar epithelial cells having inconspicuous microridges. Ultrastructurally, the mucosal surface of the intestine consists of mucous cells with electron dense granules and columnar epithelial cells having numerous microvilli, mitochondria, endoplasmic reticulum, lysosomes and Golgi body. Cellular components of the anterior and middle intestine participate in the absorption whereas the presence of enormous blood vessels and capillary net work of posterior intestine probably responsible for air breathing.

Key words: Lepidocephalichthys guntea, intestine, histoarchitecture, fine structure.

Introduction

Fishes exhibit enormous variations in their food habits and feeding specialization. The macroscopic and microscopic features of the alimentary canal along with associated structures also become modified accordingly. Among certain fish species, different parts of the digestive tract function as respiratory organ. The posterior intestine is used for gas exchange by some species belonging to families like Loricaridae, Cobitidae and Callichthyidae (Graham and Baird, 1982; McMahon and Burggren, 1987; Podkowa and Goniakowska-Witalińska, 2002). In some loaches, the digestive phase of the intestine exchanges with the respiratory phase