

B.A./B.Sc 4th Semester (Honours) Examination, 2019 (CBCS)**Subject : Mathematics****Paper : BMH4 CC10****(Ring Theory and Linear Algebra I)****Time: 3 Hours****Full Marks: 60***The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words
as far as practicable.*

[Notations and Symbols have their usual meaning.]

Group-A

Marks : 20

1. Answer any ten questions:

2×10=20

- (a) Show that a ring R is commutative if $x^3 = x$ for all $x \in R$.
- (b) What are ideals of a field? Justify your answer.
- (c) Suppose R is the ring of all real valued continuous functions defined on the closed interval $[0, 1]$ and let $S = \{f \in R : f(\frac{1}{2}) = 0\}$. Then S is an ideal of R . — Justify.
- (d) The ring $\left\{ \begin{pmatrix} a & b \\ 2b & a \end{pmatrix} : a, b \in \mathbb{R} \right\}$ is a field. — Justify.
- (e) Suppose F is a field with 2^n elements, where $n \in \mathbb{N}$. Find the characteristic of F .
- (f) Define a homomorphism from the ring \mathbb{Z} of integers into the ring \mathbb{Z}_5 of integers module 5.
- (g) Give an example to show that a quotient ring of an integral domain may not be a field.
- (h) Let R be a commutative ring of characteristic 2. Define a map $\varphi: R \rightarrow R$ by $\varphi(a) = a^2 \forall a \in R$. Prove that φ is a ring homomorphism.
- (i) Is $(0, 0, 1)$ a linear combination of $(1, 0, 1)$ and $(0, 1, 1)$? Justify your answer.
- (j) If $S = \{(1, 0, 0), (0, 1, 0)\}$, describe geometrically the linear span of S in the real vector space \mathbb{R}^3 .
- (k) Is the union of two subspaces of a vector space V a subspace of V ? Justify your answer.
- (l) Let V be a finite dimensional vector space and W be a subspace of V . What is the relation among $\dim V/W$, $\dim V$ and $\dim W$?

- (m) Is the map $T(x, y) = (x, y + 3), \forall x, y \in \mathbb{R}$ a linear transformation from the real vector space \mathbb{R}^2 into itself? Justify your answer.
- (n) State the rank-nullity theorem for vector spaces.
- (o) It is given that the map $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$ defined by $T(x, y) = (x, x + y, y) \forall x, y \in \mathbb{R}$ is a linear transformation from the real vector space \mathbb{R}^2 to the real vector space \mathbb{R}^3 . Find $\ker T$.

Group-B

Marks : 20

2. Answer any four questions:

5×4=20

- (a) (i) Let $\mathbb{Z}[\sqrt{2}] = \{a + b\sqrt{2} | a, b \in \mathbb{Z}\}$. Prove that $\mathbb{Z}[\sqrt{2}]$ is a ring under the usual addition and multiplication of real numbers.
- (ii) Let R be a ring and α be a fixed element of R . Show that $I_\alpha = \{x \in R | \alpha x = 0\}$ is a subring of R . 3+2=5
- (b) (i) Let n be a positive integer. Prove that $n\mathbb{Z}$ is a prime ideal of the ring \mathbb{Z} of integers if and only if n is prime.
- (ii) Let R be a ring with unity 1. Prove that R has characteristic $n (\neq 0)$ if and only if n is the smallest positive integer such that $n \cdot 1 = 0$. 3+2=5
- (c) (i) Let φ be a homomorphism from a ring R onto a ring S . If I is an ideal of R , prove that $\varphi(I)$ is an ideal of S .
- (ii) State the third isomorphism theorem for rings. 3+2=5
- (d) Let U and W be subspaces of a vector space V over a field F . Prove that
- (i) $U + W = \{u + w | u \in U, w \in W\}$ is a subspace of V .
- (ii) $U + W$ is the smallest subspace of V containing U and W . 3+2=5
- (e) (i) Find a basis for the real vector space \mathbb{R}^3 that contains the vectors $(1, 2, 1)$ and $(3, 6, 2)$.
- (ii) It is given that $W = \{(x, y, z) | x, y, z \in \mathbb{R}, 2x + y - z = 0\}$ is a subspace of the real vector space \mathbb{R}^3 . Find the dimension of W . 3+2=5
- (f) (i) Let U and V be two finite dimensional vector spaces over the same field F such that $\dim U = \dim V$. Show that U and V are isomorphic vector spaces.
- (ii) It is given that the map $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T(x, y) = (x + y, x - y) \forall x, y \in \mathbb{R}$ is a linear transformation from the real vector space \mathbb{R}^2 into itself. Find $\dim(\text{Im}T)$. 3+2=5

Group-C

Marks : 20

3. Answer any two questions: 10x2=20

(a) (i) Let $M_{2 \times 2}(\mathbb{Z})$ be the ring of all 2×2 matrices over the integers and let $R = \left\{ \begin{pmatrix} a & a-b \\ a-b & b \end{pmatrix} \mid a, b \in \mathbb{Z} \right\}$. Prove or disprove that R is a subring of $M_{2 \times 2}(\mathbb{Z})$.

(ii) Prove that a finite integral domain is a field.

(iii) Let $\mathbb{R}[x]$ be the ring of polynomials in x with real coefficients and $\langle x^2 + 1 \rangle$ be the principal ideal of $\mathbb{R}[x]$ generated by $x^2 + 1$. Prove that $\langle x^2 + 1 \rangle$ is a maximal ideal of $\mathbb{R}[x]$. 2+4+4=10

(b) (i) Let R be a commutative ring with unity and A be an ideal of R . Prove that R/A is a field if and only if A is maximal.

(ii) State and prove the first isomorphism theorem for rings. 5+5=10

(c) (i) Let $S = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a + b = 0, a, b, c, d \in \mathbb{R} \right\}$. Prove that S is a subspace of the vector space $M_{2 \times 2}(\mathbb{R})$ of all 2×2 real matrices. Find the dimension of S .

(ii) Show that the set of vectors $S = \{(1, 2, 0), (2, 1, 3), (1, 1, 1), (2, 3, 1)\}$ of vectors is linearly dependent in the real vector space \mathbb{R}^3 . Find a linearly independent subset T of S such that $L(T) = L(S)$. ($L(A)$ denotes the linear span of A)

(iii) It is given that $U = \{(x, y, z) \in \mathbb{R}^3 \mid x + 2y = z\}$ and $W = \{(x, y, z) \in \mathbb{R}^3 \mid 2x + 2z = y\}$ are subspaces of the real vector space \mathbb{R}^3 . Find a basis for the subspace $U \cap W$. 5+3+2=10

(d) (i) Let $a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0$

$$a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0$$

.....

.....

$$a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0$$

be a system of m linear homogenous equations with real coefficients in n variables, where $n > m$. Using Rank-nullity theorem show that the system has a non-trivial solution.

(ii) Let V be a vector space with a basis $\{e^{3t}, te^{3t}, t^2e^{3t}\}$ over the field of real numbers.

$D : V \rightarrow V$ be defined by $D(f(t)) = \frac{d}{dt}f(t) \forall f(t) \in V$. Find the matrix of D in the given basis.

(iii) Give an example of a linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that $T^2(\alpha) = \alpha \forall \alpha \in \mathbb{R}^2$. 3+5+2=10